

Background: Neural constraints


Gregor Schöner
dynamicfieldtheory.org

the brain

motor
cortex



frontal
cortex

visual
cortex



to motor
output


neurons




$\sim 10^{11}$ with 10000 synapses each

neurons


A



B




C



neurons


four components of neurons



Functional analysis of the brain

■ vary conditions “outside” the brain: stimulus, motor task, cognitive task and relate to neural activity: **coding/decoding, cognitive neuroscience**

■ or the reverse: vary neural substrate (lesioning, optogenetics, etc) and observe what happens to behavior/competence: **neuropsychology**



- Neurophysics
- Sensors, actuators, rate code
- Receptive fields, tuning curves
- Maps
- Distributions of population representation
- Patterns of connectivity
- Synaptic dynamics



Neuro-physics

- membrane potential, $u(t)$, evolves as a dynamical system

$$\tau \dot{u}(t) = -u(t) + h + \text{input}(t)$$



$\tau \approx 10$ ms time scale

- only when membrane potential exceeds a threshold is activation transmitted to downstream neurons




[from: Tresilian, 2012]

temporal summation



Neural dynamics


- replace spiking mechanism by sigmoid:
 - low levels of activation: not transmitted to downstream systems
 - high levels of activation: transmitted to downstream systems
- abstracting from biophysical details ~ **population level membrane potential**




- Neurophysics
- Sensors, actuators, rate code
- Receptive fields, tuning curves
- Maps
- Distributions of population representation
- Patterns of connectivity
- Synaptic dynamics

Sensor cells


Vision




Smell




Taste




Touch




Thermal senses




Pain




Hearing




Balance



Proprioception




Sensor characteristic




[from: Tresilian, 2012]

Sensor characteristic

A Neural code of stimulus magnitude




B Perceived sensation intensity




Sensor characteristic

photoreceptors

A Phototransduction and neural signaling




C Voltage response to light




[from: Tresilian, 2012]

Motor neurons



[from: Tresilian, 2012]

Motor neurons




[from: Tresilian, 2012]

Peripheral neural circuits

■ stretch reflex


B Monosynaptic pathways (stretch reflex)



- Neurophysics
- Sensors, actuators, rate code
- Receptive fields, tuning curves
- Maps
- Distributions of population representation
- Patterns of connectivity
- Synaptic dynamics

Sensory networks


■ retinal network




Sensory networks

■ receptive fields

(A) A ganglion cell is connected to the photoreceptors occupying a roughly circular region




(B) Receptive fields of neighboring ganglion cells overlap




Sensory networks

space-time structure of receptive fields


retinal ganglion cells



Sensory networks




 LGN





Sensory networks

■ tuning curves in primary visual cortex




tuning curve: sensory system

■ Hubel, Wiesel: primary visual cortex (monkey)



tuning curve: motor system


■ primary motor cortex (monkey)





- Neurophysics
- Sensors, actuators, rate code
- Receptive fields, tuning curves
- Maps
- Distributions of population representation
- Patterns of connectivity
- Synaptic dynamics

Neural maps



- tuning curves studied systematically across the cortical surface
- => feature maps
- topography




B Ocular dominance columns



C Orientation columns




D Blobs, interblobs (V1), and stripes (V2)




Thin stripe

Thick stripe



[Charles D. Gilbert,
Aniruddha Das, Chapter 21
of Kandel et al 2021]


Motor networks



- Neurophysics
- Sensors, actuators, rate code
- Receptive fields, tuning curves
- Maps
- Distributions of population representation
- Patterns of connectivity
- Synaptic dynamics

Population code


- notion that all activated neurons contribute to feature representation according to their tuning curves



[Charles D. Gilbert, Aniruddha Das, Chapter 21 of Kandel et al 2021]

Experimental evidence for population representations

- Lee, Rohrer, Sparks: use the topographic map of saccadic endpoint in superior colliculus
- to reversibly deactivate portions of the population: observe predicted deviations of saccadic endpoints



[after Lee, Rohrer, Sparks: Nature (1988)
in Chapter 3 of the book]


Population representation in the visual system

- Example 1: Jancke et al: A17 in the cat, population representation of retinal location

Jancke, Erlhagen, Dinse, Akhavan, Giese, Steinhage,
Schöner JNeurosci 19:9016 (99)

Distribution of Population Activation (DPA)


- determine tuning to retinal location for each cell
- superpose tuning curves weighted by current firing rate: **distribution of population activation DPA** representing retinal location



Neural grounding of DFT: sensory

■ DPA of stimuli presented to all neurons

elementary stimuli




2.8°

nasal

temporal

current stimulus:
square of light

range of retinal field
sampled by neurons




□ 0.4°




Neural grounding of DFT: sensory

■ => observe interaction in DPA




interaction




activation level in DPA




at location of left component stimulus




response to composite stimuli



superposition of responses to each elemental stimulus




evidence for inhibitory interaction



time


[ms]

model by dynamic field:






Neural grounding of DFT: sensory

- DPA of orientation and (ID) retinal location




Population representation motor system

- motor and pre-motor cortex (macaque)
- in behaving animal





Complete Information




Distribution of Population Activation (DPA)

Distribution of population activation = $\sum_{\text{neurons}} \text{tuning curve} * \text{current firing rate}$



[Bastian, Riehle, Schöner, 2003]

Decision making in DPA



[Cisek, Kalaska 2005]


Distributions of Population Activation (DPA)

- neurons are not localized within DPA!
- cortical neurons really are sensitive to many dimensions
 - motor: arm configuration, force direction
 - visual: many feature dimensions such as spatial frequency, orientation, direction...
- => DPA is a projection from that high-dimensional space onto a single dimension


- Neurophysics
- Sensors, actuators, rate code
- Receptive fields, tuning curves
- Maps
- Distributions of population representation
- Patterns of connectivity
- Synaptic dynamics

Forward connectivity


(A) Disynaptic connectivity between P and Q



(B) Convergent pathway






(C) Divergent pathway



Dale's law

- all synaptic connections coming from a given neuron are of the same type



Patterns of connectivity

■ recurrent connectivity




- Neurophysics
- Sensors, actuators, rate code
- Receptive fields, tuning curves
- Maps
- Distributions of population representation
- Patterns of connectivity
- Synaptic dynamics

Learning by synaptic plasticity

- synaptic strengths change as a function of pre/post synaptic neural state

Hebbian Learning in experiments (schematic)





Learning by plasticity

- spike-time dependent plasticity

- strengthening of synapses in which pre-synaptic spike precedes post-synaptic spike

- weakening synapses when the temporal order is the reverse...

Spike-time dependent plasticity



Pre
before post

[Gerstner et al, 2014]

- Neurophysics
- Sensors, actuators, rate code
- Receptive fields, tuning curves
- Maps
- Distributions of population representation
- Patterns of connectivity
- Synaptic dynamics